

GUIDE TO MARINE INVADERS IN RI COASTAL WATERS

Undaria pinnatifida undaria kelp, wakame seaweed

Potential
Invader



Photos: Steve Lonhart /Monterey Bay NMS



PHYSICAL DESCRIPTION


Brown kelp with wide, brown to yellow-tan blades

Mature plants have a divided blade (with pinnae) with conspicuous midrib, holdfast, stipe (stem), and spiral, folded sporophyll

Young plants have holdfast, stipe, undivided blade, and initially no midrib

Grows rapidly to 3-9 ft (1-3 m) long

Undaria pinnatifida
young plant



Undaria pinnatifida
mature plant
(not to scale)




Illustration: © Rob Gough, DFBW
Photo: Steve Lonhart /MBNMS

HABITAT PREFERENCE

- Found from low intertidal to subtidal depths of approx. 15 ft (5 m)
- Grows on hard surfaces including rocks, ropes, docks, pilings, moorings, and other structures
- Can form dense 'kelp forests' in sheltered waters

1

2

3

4

5

6

7

8

GUIDE TO MARINE INVADERS IN RI COASTAL WATERS

Undaria pinnatifida undaria kelp, wakame seaweed

Potential
Invader

INVASION STATUS & ECOLOGICAL CONCERNS

Undaria pinnatifida is an invasive kelp native to Japan. Commonly known as 'wakame', an ingredient in miso soup, this alga is commercially grown throughout Asia for human consumption. It has invaded many of the world's oceans, from European waters, to waters off New Zealand. In 2001, *U. pinnatifida* was discovered on North American shores in Monterey Bay, CA, where efforts to manually remove it continue today.

This alga is a very prolific and hardy species, with a growth rate measured at 1-2 cm per day, and a max. length of 9 ft (3 m). It can establish itself on a wide variety of surfaces. As a result of its amazing growth rate and wide blades, this seaweed quickly starves the natural understory algae of light. As these algae die, fish and invertebrates must move elsewhere to find food, thus creating extra feeding pressure on adjacent areas. *U. pinnatifida* has drastically affected the ecosystem of many waters that it has invaded. In addition to such ecological damages, it is a fouling species on ship hulls, nets, fishing gear, moorings, ropes, and other structures, and as such, increases labor and maintenance costs.

SIMILAR SPECIES

Saccharina latissima (sugar kelp)

Undaria pinnatifida may be mistaken for our region's native brown kelps, particularly *Saccharina latissima*. However, *S. latissima* has ruffled edges along undivided blades with no midrib, vs. divided blades and conspicuous midrib of *U. pinnatifida*.

While several of the Gulf of Maine's native kelp species possess some of the features of *U. pinnatifida* (i.e. midrib in *Alaria* sp., divided blades of *Laminaria digitata*), none possess the folded, spiral sporophyll.

(left to right):

S. latissima blade, stipe and holdfast
S. latissima blade showing absence of midrib

Alaria sp. showing midrib



This card is adapted from an original series produced by Salem Sound Coastwatch (www.salemsound.org). The original series was funded by the MA EOEPA Office of Coastal Zone Management with funding from the U.S.F.W.S. For more information please visit www.mass.gov/czm/invasives/monitor/reporting.htm. The production of this adapted card was funded by the RI Coastal Resources Management Council with funding from the U.S. Fish & Wildlife Service. To report findings please email kcute@crmc.ri.gov or call (401) 783-7772 or (401) 783-3370.