



Oliver Stedman Government Center
4808 Tower Hill Road, Suite 116
Wakefield, RI 02879
401-783-3370

**PUBLIC NOTICE for PROPOSED RULE-MAKING and
PUBLIC HEARING**

**Proposed Amendments to the Coastal Resources Management Program
Section 145 – Climate Change and Sea Level Rise**

Pursuant to Chapter 46-23 of the State of Rhode Island General Laws, as amended, the Coastal Resources Management Council proposes to amend and take public comment on the following sections only of the Coastal Resources Management Program (CRMP): **Section 145 – Climate Change and Sea Level Rise**. Comments will not be taken on any other section of the Coastal Resources Management Program during this notice period. In accordance with the procedures of the RI Administrative Procedures Act (R.I.G.L. § 42-35-3) and the Rules and Regulations of the Coastal Resources Management Council, notice is hereby given of the intent of the Coastal Resources Management Council to hold a public hearing, accept public comment, and afford interested persons reasonable opportunity to submit data, views or arguments orally or in writing during the 30-day comment period and the public hearing.

The **public hearing** will be held at **6:00 p.m. on January 26, 2016 in Conference Room A at the Department of Administration, One Capitol Hill, Providence, RI**. The room is accessible to the disabled and persons requesting interpreter services for the hearing impaired must notify the Council office at 401-783-3370 or RI 711 at least three (3) business days in advance of the hearing date so that such assistance can be provided at no cost to the person requesting.

Summary of Proposed Amendment to CRMP Section 145:

The purpose of the proposed amendment to Section 145 of the Coastal Resources Management Program is to update sea level rise (SLR) projections for short-, mid- and long-term timelines of 2035, 2050 and 2100, respectively, as calculated using current National Oceanic Atmospheric Administration (NOAA) methodology and based on the Newport, RI NOAA tide gauge. The CRMC's existing policy in Section 145 is to periodically review new scientific information regarding SLR. The proposed amendment would provide the state and municipalities with appropriate sea level rise projections for consideration in the planning and design of future public and private investments and natural resource restoration projects.


The Council has complied with the requirements of R.I. General Laws §§ 42-35.1-3 and 42-35.1-4 and has filed copies of the proposed regulations with the Governors Office and the Office of Regulatory Reform of the Department of Administration's Office of Management and Budget. The Council has determined that the proposed regulations do not overlap or duplicate any other state regulation and will not have any adverse effect on small businesses.

The proposed regulations are available on the CRMC website – www.crmc.ri.gov. Additionally, the proposed regulations can be reviewed in person at the Coastal Resources

Management Council offices located at the Oliver Stedman Government Center, 4808 Tower Hill Road, Wakefield, RI. Further information may be obtained by contacting the Coastal Resources Management Council offices at 783-3370.

All interested persons are invited to submit written comments on the proposed amendments to Section 145 of the Coastal Resources Management Program by the close of business on **January 12, 2016** to provide advance notice to the Council prior to the public hearing. All such comments should be directed to Grover J. Fugate, Executive Director, at the above address.

Signed this **11th** day of **December**, 2015



Jeffrey M. Willis, Deputy Director
Coastal Resources Management Council

Proposed Amendment

RI Coastal Resources Management Program – Section 145 Climate Change and Sea Level Rise

Revise existing Section 145 as follows:

Unaltered text is existing text in Section 145, while new text is **underlined** and deleted text is **~~struck-through~~**

Section 145 Climate Change and Sea Level Rise

A. Definitions

1. Climate is the long-term weather average observed within a geographic region, and climate change refers to fluctuations in the Earth's climate system as a result of both natural and anthropogenic causes. Currently the long term climate change trend is evidenced by rising global temperatures; increasing extremes within the hydrologic cycle resulting in more frequent floods and droughts; and rising sea level.

2. Sea level is the height of the sea with respect to a horizontal control point or benchmark such as the National Geodetic Vertical Datum of 1929 (NGVD 29) or the North American Vertical Datum of 1988 (NAVD 88). Sea level rise refers to the net increase in mean sea level over time in response to global climate, local tectonic changes, glacial isostatic adjustment, and ocean dynamics.. Sea level rise indicates a positive trend, thus an increase in sea level as compared to historic measurements. Global sea level rise is the worldwide variations in sea level due to eustatic contributions such as thermal expansion of seawater and melting glacial ice sheets. Relative sea level rise is a regional change in sea level relative to land surface elevations.

Relative sea level rise is influenced by tectonic response to ice or sediment loading, land subsidence due to extraction of water or oil, dynamic effects of ocean currents or the gravitational pull of ice sheets on ocean waters. Sea levels are rising along most of the world's coastlines, including Rhode Island. However, in places that are experiencing rapid uplift due to tectonic plate movement or glacial isostatic adjustment the relative sea level trends are falling because the land is rising faster than the sea (<http://tidesandcurrents.noaa.gov/sltrends/sltrends.shtml>).

3. Vertical datums are either fixed benchmarks such as NGVD 29 and NAVD 88 or site specific tidal datums such as mean high water, mean low water and mean sea level. NGVD 29 is based on the local mean sea level in 1929, which has changed over time. NAVD 88 is now the official civilian vertical datum for surveying and mapping activities in the United States. The conversion to NAVD 88 should be accomplished on a project-by-project basis. It should be noted, however, that NAVD 88 is not synonymous with mean sea level nor does it correct for sea level changes that have occurred since the establishment of NGVD 29. Tidal datums, such as mean sea level (MSL) or mean high water (MHW) vary according to the specific location, and represent the mean heights observed over the National Tidal Datum Epoch. Datum conversions for NOAA tide gauges at www.tidesandcurrents.noaa.gov or can be calculated using the NOAA VDatum software available at <http://vdatum.noaa.gov> or the US Army Corps of Engineers CORPSCON software available at <http://www.agc.army.mil/corpscon>

4. Horizontal datums are either fixed benchmarks or site-specific control points that establish location for a point on a map consistent with a coordinate system. The North American Datum of 1983 (NAD 83) is the official horizontal datum for the United States based on a geocentric origin and the Geodetic Reference System 1980. Conversions between the datums can be made using software such as NADCON available from the National Geodetic Survey at http://www.ngs.noaa.gov/PC_PROD/pc_prod.shtml or the US Army Corps of Engineers CORPSCON software at <http://www.agc.army.mil/corpscon/>.

B. Findings

1. On very long (geologic) time scales, sea level naturally fluctuates in response to variations in astronomical configurations that cause changes in the Earth's climate system. Since the Last Glacial Maximum (approximately 26,000 years ago), global sea level has risen by over 390 feet (120 meters), as water that was previously trapped in continental ice sheets has made its way into the global ocean.

2. Sea level rise is a direct consequence of global climate change. Greenhouse gas emissions to the atmosphere increase surface warming, which in turn increases the volume of ocean waters due to thermal expansion, and accelerates the melting of glacial ice. Atmospheric greenhouse gas concentrations are already higher than levels at the last interglacial period, when sea levels were 13 to 19 feet (4 to 6 meters) higher than at present (Overpeck et al., 2006). Greenhouse gas concentrations are expected to continue to increase through 2100.


3. Human activities and increased concentrations of greenhouse gasses in the atmosphere have accelerated the historic rate of eustatic sea level rise. Over the last 100 years, sea levels have risen 0.56 feet (0.17 m) globally. The average rate of rise during the years between 1961 and 2003 was 0.071 inches per year (1.8 mm/yr), and between 1993 and 2003 the rate nearly

doubled to 0.12 inches per year (3.1 mm/yr) (IPCC, 2007). [The present rate of global sea level rise is 3.3 mm/yr as measured by satellite altimetry. See: http://sealevel.colorado.edu/.](http://sealevel.colorado.edu/)

4. In addition to rising global sea levels, the land surface in Rhode Island was believed to be subsiding at a rate of approximately 6 inches (15 cm) per century (Douglas, 1991). More recent studies indicate that many more factors, including changes in ocean circulation, contribute to Rhode Island's relative sea level rise than subsidence alone. The combination of these effects is evident from the long-term trend recorded by the Newport tide gauge (Figure 1), which indicates a rate of 10.68 inches (26.927.4 cm) of relative sea level rise per century or 2.6974 mm per year.

5. The rate of sea level rise is accelerating. Future sea level rise, like the recent rise, is not expected to be globally uniform or linear. Some regions will become more substantially inundated than the global average, and others less. Of foremost concern is the trend in eustatic rise as observed from tide-gauge records over the past century. The rate of rise globally during the past 20 years is 25% faster than the rate of rise in any 20 year period that exists in the instrumental record (Church and White, 2006; Rahmstorf et al., 2007, Vermeer and Rahmstorf, 2009 and Rahmstorf et al., 2011).

6. Model-simulated projections of global sea level over the 21st century also clearly demonstrate accelerated progression. Predictions have ranged from 4 inches (10 cm) to several feet above current levels by the year 2100. As a rule, sea level estimates are increasing as the science of modeling becomes more developed.



Adapted from: http://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?stnid=8452660%20Newport,%20RI



Boothroyd 2012

The above graphic will be deleted and replaced with the graphic below




Figure 1. Sea level has risen ~~8.59~~1 inches since ~~1931~~1930 based on the long-term trend at Newport.

7. When compared with actual observations, modeling scenarios can be quite conservative, as recently observed rates of continental ice melt are greater than those used to generate estimates of sea level rise over the coming century. Since 1990, sea level has been rising faster than the rate predicted by models used to generate IPCC (2001) estimates (Rahmstorf et al., 2007).

8. Higher global temperatures indicate a greater risk of destabilizing the Greenland and West Antarctic ice sheets, yet a great amount of uncertainty remains as to the overall contribution from ice sheet melting. The recent and much publicized ~~Fourth~~Fifth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC ~~2007~~2013) projects ~~7-11~~ to ~~23-39~~ in (~~18 to 59~~28 to 98 cm) of eustatic sea level rise in the coming century. Sea levels are rising faster now than in the previous two millennia, and the rise is projected to accelerate – regardless of the emissions scenario, even with strong climate mitigation (IPCC, 2013). These estimates ~~do not~~ include limited contributions of ice flow dynamics and do not include local subsidence.

9. Rahmstorf (2007) and Rahmstorf et al. (2011) correlate global sea level rise to global mean surface temperature, which is a good approximation for observations of the 20th century. When this relationship is applied to 21st century warming scenarios, eustatic rise is projected between 1.6 to 4.6 feet (50 to 140 cm) above 1990 levels. Accounting for regional isostatic effects, this estimate suggests that by 2100 sea level in Rhode Island could rise approximately 2 to 5 feet (65 to 155 cm).

10. More recent scientific observations and refined climate models support previous projections and indicate that globally a range of sea level rise of between 2 to 6 feet (0.6 to 1.9 m) above 1990 levels is expected by the year 2100 (Jevrejeva et al., 2010; Vermeer and Rahmstorf, 2009 and Rahmstorf et al., 2011).

11. Regional rates of sea level rise will differ across the globe. The dynamic effects of ocean currents and the diminishing gravitational pull of dwindling ice sheets on ocean waters, have the potential to increase sea level rise rates at a particular location. Model projections indicate that a

slowdown in the Atlantic Meridional Overturning Circulation (AMOC) may lead to a rapid rise in sea level on the northeast coast of the United States (Yin et al., 2009, Yin et al., 2011, Kuhlbrodt et al., 2009, Hu et al., 2009, Bingham and Hughes, 2009 and Kopp et al., 2010). Changes in static equilibrium of ocean and ice mass distribution will have an impact on relative sea levels depending on the rate of melt (Kopp et al., 2010).

12. U.S. Geological Survey scientists detail in their study (Sallenger et al., 2012) that recently accelerated sea level rise along the Atlantic Coast will result in sea levels 8 to 11 inches (20-29 cm) higher than the global average from Cape Hatteras, NC to Boston, MA by 2100. They present evidence that the rate of sea level rise increase in the study area was 3-4 times higher than the global average during the last two tidal epochs of 1950-1979 and 1980-2009. Sea level rise combined with storm surge, wave run-up and set-up will increase the vulnerability of near-shore areas to flooding, beach erosion and coastal wetland degradation.

13. A study by Strauss et al. (2012) examines topographic vulnerability of low-lying coastal land in the continental United States to sea level rise and flooding. The researchers found that there are presently 2705 housing units along the Rhode Island shoreline that are located less than 1 meter (39 inches) above local mean high water (MHW). These housing units are most at risk for increased flooding and eventual submersion as a result of sea level rise.

14. Tibaldi et al. (2012) investigated the historic patterns of extreme high tide events at 55 coastal locations of the contiguous United States using a detailed analysis of the NOAA tide gauge station data from 1979-2008 coupled with anticipated relative sea level rise. They calculate an increase of 5.1 inches (0.13m) by 2030 and 12.2 inches (0.31m) by 2050 above the 2008 mean high water level as measured at the Newport tide gauge. The study indicates that the frequency of extreme high tide levels will increase significantly in the coming years.

15. Climate change will result in wide scale systematic changes in the terrestrial and marine environments. These changes will result in ecosystem shifts that will challenge natural resource managers' efforts to cope and adapt to the new regime.

16. Future increases in relative sea level will displace coastal populations, threaten infrastructure, intensify coastal flooding and ultimately lead to the loss of recreation areas, public space, and coastal wetlands.

17. Coastal infrastructure will become increasingly susceptible to complications from rising sea levels, as the upward trend continues. Residential and commercial structures, roads, and bridges will be more prone to flooding. Sea level rise will also reduce the effectiveness and integrity of existing seawalls and revetments, designed for historically lower water levels.

18. Higher sea levels will result in changes in surface water and groundwater characteristics. Salt intrusion into aquifers will contaminate drinking water supplies and higher water tables will compromise wastewater treatment systems in the coastal zone.

19. Future increase in relative sea level will increase the extent of flood damage over time. Lower elevations will become increasingly susceptible to flooding as storm surge reaches further inland due to sea level rise in concert with a probable increase in the intensity of storms predicted from climate change. As a result, more coastal lands will be susceptible to erosion.

20. At historic rates of sea level rise, the relative surface elevation of a salt marsh may be maintained through the process of accretion (the build-up of live and decaying plant parts and inorganic sediments). Yet, at high rates of relative sea level rise as predicted by Rahmstorf (2007), accretive processes in coastal wetlands will not keep pace. These habitats can become submerged resulting in a loss of salt marsh vegetation and an alteration of habitat types. This has been demonstrated by the rapid salt marsh loss in coastal Louisiana. Observations by environmental researchers here in Rhode Island indicate that salt marshes are losing high marsh habitat as a result of more frequent inundation and possibly a consequence of accretion rates that are unable to keep pace with increased rates of sea level rise. As salt marshes and other coastal habitats become submerged, they migrate inland. However, coastal development has decreased the amount of upland open space adjacent to these habitats limiting their ability to migrate landward. Thus, an increase in the rate of relative sea level rise will likely result in significant losses of coastal saltmarsh habitats.

21. The average annual temperature of southern New England coastal waters, including Narragansett Bay, has risen approximately two (2) degrees Fahrenheit since the 1960's. This warming trend is implicated in the change of species composition and abundance in Narragansett Bay waters (Nixon, et al., 2003).

22. Increased water temperatures due to climate change will work synergistically with high nutrient levels to stress eelgrass beds. Eelgrass grows best in cool, clean waters. Even as nutrient levels in the Bay are reduced from wastewater treatment plants, if Bay and coastal waters continue to warm due to climate change, it will adversely impact eelgrass beds (Bintz, et al., 2003).

23. Barrier islands are forced landward with rising sea levels. Increased frontal erosion and retreat of the barriers will cause Rhode Island's south shore to migrate continuously landward with rising sea levels.

24. Due to the timescales associated with climate processes and feedbacks, anthropogenic warming and sea level rise will continue for centuries regardless of steps taken to curb greenhouse gas emissions (IPCC, 2007).

25. Flooding is a destructive natural hazard and results in economic loss to the citizens of Rhode Island. Approximately 154 square miles (14%) of the State's 1100 square miles of land area are mapped as Special Flood Hazard Areas by the National Flood Insurance Program (NFIP) where there is a 1% chance of flooding in any given year. (RIEMA, 2011). More than 16,000 buildings are located within these flood prone areas with an additional 12,000 buildings located in areas mapped as 0.2% chance of flooding (based on CRMC GIS assessment of E911 data and flood zones).

26. All 39 communities within the State participate in the National Flood Insurance Program, yet only about half of Rhode Island property owners located within Special Flood Hazard Areas carry flood insurance (RIEMA and E911 data assessment).

27. Pursuant to R.I.G.L. § 46-23-6, the Council is authorized to develop and adopt policies and regulations necessary to manage the coastal resources of the state and protect life and

property from coastal hazards resulting from projected sea level rise and probable increased frequency and intensity of coastal storms due to climate change. The Council is also authorized to collaborate with the State Building Commissioner and adopt freeboard calculations (a factor of added safety above the anticipated flood level), in accordance with R.I.G.L. § 23-27.3-100.1.5.5.

28. The U.S. Army Corps of Engineers (USACE) ~~has~~ released a revised circular dated ~~October 1, 2011~~[December 31, 2013](#) detailing its methodologies for assessing the impacts of sea level rise in the planning, design, engineering, construction, operation and maintenance of USACE civil works projects in coastal areas. The required project analyses determine how sea level rise scenarios may affect risk levels to the surrounding area and identify the design or operations and maintenance measures that will minimize adverse consequences while maximizing the beneficial effects of the project. See: <http://publications.usace.army.mil/publications/eng-circulars/EC-1165-2-212.pdf> http://www.publications.usace.army.mil/Portals/76/Publications/EngineerRegulations/ER_1100-2-8162.pdf. In addition, the USACE in collaboration with the National Oceanic and Atmospheric Administration (NOAA) have released a sea level rise calculator available online at: <http://corpsclimate.us/ccaceslcurves.cfm>. The two NOAA tide gauges applicable to Rhode Island when using the sea level rise calculator are located in Providence and Newport.

29. NOAA has very high confidence that global mean sea level will rise at least 0.2m (8 inches) and no more than 2.0m (6.6 feet) by 2100 (Parris et al., 2012).

~~29~~30. According to a USGS report (Titus et al., 2009), preparing in advance for expected sea level rise is justifiable for several types of impacts, as it may be less costly to react now than to react to an adverse condition in the future. Some examples:

- Coastal wetland protection. Preserving undeveloped lands abutting coastal wetlands allows wetland migration, but once developed, it is very difficult to make land available for wetland migration. Therefore, it is far more practicable to promote wetland migration by setting aside land before it is developed and preserving coastal buffer zones, than to require development to be removed as sea level rises.
- Some long-term infrastructure. Whether it is beneficial to design coastal infrastructure to anticipate rising sea level depends on economic analysis of the incremental cost of designing for a higher sea level now, and the retrofit cost of modifying the structure at some point in the future. Most long-lived infrastructure in the threatened areas is sufficiently sensitive to rising sea level to warrant at least an assessment of the costs and benefits of preparing for rising sea level.
- Floodplain management. Rising sea level increases the potential disparity between rates and risk. Even without considering the possibility of accelerated sea level rise, the National Academy of Sciences and a Federal Emergency Management Agency (FEMA)-supported study by the Heinz Center recommended to Congress that insurance rates should reflect the changing risks resulting from coastal erosion.

C. Policies

1. The Council will review its policies, plans and regulations to proactively plan for and adapt to climate change and sea level rise. The Council will integrate climate change and sea level rise scenarios into its programs to prepare Rhode Island for these new, evolving conditions

and make our coastal areas more resilient.

2. The Council's sea level rise policies are based upon the CRMC's legislative mandate to preserve, protect, and where possible, restore the coastal resources of the state through comprehensive and coordinated long-range planning.

3. The Council recognizes that sea level rise is ongoing and its foremost concern is the accelerated rate of rise and the associated risks to Rhode Island coastal areas today and in the future. ~~Accordingly, for planning and management purposes, it is the Council's policy to accommodate a base rate of expected 3 to 5 foot rise in sea level by 2100 in the siting, design, and implementation of public and private coastal activities and to insure proactive stewardship of coastal ecosystems under these changing conditions. It should be noted that the 3-5 ft. rate of sea level rise assumption embedded in this policy is relatively narrow and low.~~ The Council recognizes that the lower the sea level rise estimate used, the greater the risk that policies and efforts to adapt sea level rise and climate change will prove to be inadequate. Therefore, the policies of the Council may take into account different risk tolerances for differing types of public and private coastal activities. In addition, ~~this long-term sea level change base rate will be revisited by~~ the Council will regularly review periodically to address new scientific evidence regarding sea level change.

4. The Council relies upon the most recent NOAA sea level rise data to address both short- and long-term planning horizons and the design life considerations for public and private infrastructure. The Council's policy is to adopt and use the sea level change scenarios published by NOAA in Technical Report OAR CPO-1 (Parris et al., 2012), and the sea level rise change curves for Newport and Providence as provided in the USACE sea level rise calculator. As of 2015 the range in sea level rise change is projected by NOAA to be a maximum of approximately 1.0 foot in 2035, 2.0 feet in 2050 and 7.0 feet in 2100. In addition, the Council adopts and recommends the use of the StormTools online mapping tool developed on behalf of the CRMC by the University of Rhode Island Ocean Engineering program to evaluate the flood extent and inundation from sea level rise and storm surge.

D. References

Bingham, R. J. and C. W. Hughes. 2009. Signature of the Atlantic meridional overturning circulation in sea level along the east coast of North America, *Geophysical Research Letters*, Vol 36, L02603, 5 pp., doi:10.1029/2008GL036215

Bintz, J., S. Nixon, B. Buckley, and S. Granger. 2003. Impacts of temperature and nutrients on coastal lagoon plant communities. *Estuaries* Vol. 26, No. 3, p. 765-776.

Church, J.A., & White, N.J. (2006). A 20th century acceleration in global sea-level rise. *Geophysical Research Letters*, 33, L01602-L01604.

Douglas, B.C. (1991). Global sea level rise. *Journal of Geophysical Research*, 96(C4), 6981-6992.

Hu, A., G. A. Meehl, W. Han, and J. Yin. 2009. Transient response of the MOC and climate to potential melting of the Greenland Ice Sheet in the 21st century. *Geophysical Research Letters*,

Vol 36, L10707, 6 pp., doi:10.1029/2009GL037998

IPCC. (2001). *Climate Change 2001: The Scientific Basis. Contribution of Working Group 1 to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK and New York, NY: Cambridge University Press.

[IPCC. \(2007\). Climate Change 2007: The Physical Science Basis. Summary for Policymakers. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Geneva, Switzerland: UNEP.](#)

[IPCC. \(2013\) *Climate Change 2013: The Physical Science Basis. Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Geneva, Switzerland: UNEP.](#)

Jevrejeva, S., J.C. Moore, and A. Grinsted. 2010. How will sea level respond to changes in natural and anthropogenic forcings by 2100? *Geophysical Research Letters*, Vol. 37, L07703

Kopp, R. E. J. X. Mitrovica, S. M. Griffies, J. Yin, C. C. Hay, and R. J. Stouffer. 2010. The impact of Greenland melt on local sea levels: a partially coupled analysis of dynamic and static equilibrium effects in idealized water-hosing experiments. *Climatic Change* (2010) 103:619–625, DOI 10.1007/s10584-010-9935-1

Kuhlbrodt, T., S. Rahmstorf, K. Zickfeld, F. B. Vikebø, S. Sundby, M. Hofmann, P. M. Link, A. Bondeau, W. Cramer, and C. Jaeger. 2009. An Integrated Assessment of changes in the thermohaline circulation. *Climatic Change* (2009) 96:489–537, DOI 10.1007/s10584-009-9561-y

Nixon, S., S. Granger, and B. Buckley. 2003. *The Warming of Narragansett Bay. 41° North*. Vol. 2, Issue 1. Rhode Island Sea Grant and the University of Rhode Island Coastal Institute.

Overpeck, J.T., Otto-Bliesner, B.L., Miller, G.H., Muhs, D.R., Alley, R.B., & Kiehl, J.T. (2006). Paleoclimate evidence for future ice-sheet instability and rapid sea-level rise. *Science*, 311, 1747-1750.

[Parris, A., P. Bromirski, V. Burkett, D. Cayan, M. Culver, J. Hall, R. Horton, K. Knuuti, R. Moss, J. Obeysekera, A. Sallenger, and J. Weiss. 2012. *Global Sea Level Rise Scenarios for the U.S. National Climate Assessment*. NOAA Technical Report OAR CPO-1. 37 pp.](#)

Rahmstorf, S. (2007). A semi-empirical approach to projecting future sea-level rise. *Science*, 315, 368-370.

Rahmstorf, S., Cazenave, A., Church, J.A., Hansen, J.E., Keeling, R.F., Parker, D.E., & Somerville, R.C.J. (2007). Recent climate observations compared to projections. *Science*, 316(5825), 709.

Rahmstorf, S, M. Perrette, and M. Vermeer, 2011. Testing the robustness of semi-empirical sea level projections. *Climate Dynamics* DOI 10.1007/s00382-011-1226-7.

Rhode Island Emergency Management Agency (RIEMA). 2011. Rhode Island State Hazard

Mitigation Plan, April 2011.

Rhode Island State Building Code. SBC-1 and SBC-2. Department of Administration, Providence, RI. <http://www.ribcc.ri.gov/>

Sallenger, A., K. Doran and P. Howd. 2012. Hotspot of accelerated sea-level rise on the Atlantic coast of North America. *Nature Climate Change Letters*. pub. online 24 June 2012.

Strauss, B., R. Ziemiński, J. Weiss and J. Overpeck. 2012. Tidally adjusted estimates of topographic vulnerability to sea level rise and flooding for the contiguous United States. *Environmental Research Letters* 7 (2012) 014033 12pp.

Tebaldi C. , B. H Strauss and C. E. Zervas. 2012. Modeling sea level rise impacts on storm surges along US coasts. *Environmental Research Letters*, 7 (2012) 014032 (11pp)
doi:10.1088/1748-9326/7/1/014032

Titus, James G., K. Eric Anderson, Donald R. Cahoon, Dean B. Gesch, Stephen K. Gill, Benjamin T. Gutierrez, E. Robert Thieler, and S. Jeffrey Williams. 2009. Coastal Sensitivity to Sea Level Rise: A Focus on the mid-Atlantic Region. Synthesis and Assessment Product 4.1. Report by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research. Washington, D.C.
ISBN 978-0-16-083086-0

Vermeer, M. and S. Rahmstorf (2009) Global sea level linked to global temperature. *Proc. Natl. Acad. Sci. USA* 106: 21527-21532

Yin J., M. E. Schlesinger, and R. J. Stouffer. 2009. Model projections of rapid sea-level rise on the northeast coast of the United States. *Nature Geoscience*, Vol. 2, DOI: 10.1038/NGEO462

[Yin J.](#), [J. T. Overpeck](#), [S. M. Griffies](#), [A. Hu](#), [J. L. Russell](#), and [R. J. Stouffer](#), 2011. Different magnitudes of projected subsurface ocean warming around Greenland and Antarctica. *Nature Geoscience* 4, 524–528 (2011) doi:10.1038/ngeo1189